

---

## NEW MEXICO HEALTH ALERT NETWORK (HAN) ADVISORY

### Acute Hepatitis A Virus Infections in Persons Experiencing Homelessness and Those Who Use Injection and Non-Injection Drugs in Bernalillo and Santa Fe Counties

05/08/2019

The New Mexico Department of Health (NMDOH) has confirmed more than 100 acute hepatitis A virus (HAV) infections, and 2 associated deaths in Bernalillo County. A confirmed case has now been identified in Santa Fe County. The outbreak has primarily affected people experiencing homelessness and persons using both injection and non-injection drugs. The Centers for Disease Control and Prevention (CDC) and state health departments are investigating hepatitis A outbreaks in multiple states among persons reporting drug use and/or homelessness and their contacts. To avoid a large-scale outbreak in our communities the NMDOH is issuing this HAN to advise clinicians.

#### Background

Since the hepatitis A outbreaks associated with person to person transmission were first identified in 2016, more than 15,000 cases, 8500 hospitalizations (57%) and 140 deaths as a result of hepatitis A infection have been reported.

Transmission in the United States is typically person-to-person through the fecal-oral route (i.e., ingestion of something that has been contaminated with the feces of an infected person). Symptoms include fever, fatigue, loss of appetite, nausea, vomiting, abdominal pain, dark urine, clay-colored bowel movements, joint pain, and jaundice. Although rare, atypical extra hepatic manifestations include rash, pancreatitis, renal disease, arthritis, and anemia. Average incubation of HAV is 28 days, but illness can occur up to 50 days after exposure. An HAV-infected person can be viremic up to six weeks and excrete virus in stool for up to two weeks prior to becoming symptomatic, making identifying exposures particularly difficult. Illness from hepatitis A is typically acute and self-limited; however, when this disease affects populations with already poor health (e.g., hepatitis B and C infections, chronic liver disease), infection can lead to serious outcomes, including death.

Person-to-person transmission of HAV between persons who report drug use and/or homelessness result from generally poor sanitary conditions and the sharing of syringes, pipes, food or drinks. Transience, economic instability, limited access to healthcare, distrust of public officials and public messages, and frequent lack of follow-up makes this population difficult to reach for preventive services such as vaccination, case management and contact tracing. These challenges make outbreaks among these groups difficult to control.

#### EPIDEMIOLOGY AND RESPONSE

1190 St. Francis Drive, N1320 • Santa Fe, New Mexico • 87502-6110  
(505) 827-0006 • FAX: (505) 827-2110 • [www.nmhealth.org](http://www.nmhealth.org)



**The best way to prevent hepatitis A infection is through vaccination with the hepatitis A vaccine.**

NMDOH has several recommendations for clinicians:

- Consider hepatitis A as a diagnosis in anyone with jaundice and clinically compatible symptoms mentioned above.
- Vaccinate the following high-risk groups:
  - Persons experiencing homelessness or transient housing
  - Persons who use injection and non-injection drugs
  - Anyone with ongoing contact with people who use injection and non-injection drugs or people who are homeless
  - Persons with chronic liver disease
  - Men who have sex with men (MSM)
  - People with direct contact with others who have hepatitis A
  
- Persons who have been exposed recently to HAV and who have not been vaccinated should receive one dose of single-antigen hepatitis A vaccine or immune globulin (IG) as soon as possible, **within 2 weeks after exposure**. Guidelines vary by age and health status (please see <https://www.cdc.gov/hepatitis/outbreaks/InterimOutbreakGuidance-HAV-VaccineAdmin.htm> for additional information).
  
- Ensure all suspected and confirmed cases of hepatitis A are reported to the NMDOH Epidemiology and Response Division within 24 hours.

To report cases or for more information please call the NMDOH Epidemiology and Response Division at 505-827-0006. For more information about hepatitis A outbreaks please see the links below.

- Use of Hepatitis A Vaccine for Persons Experiencing Homelessness  
<https://www.cdc.gov/mmwr/volumes/68/wr/mm6806a6.htm>
- [2017 – Outbreaks of hepatitis A in multiple states among people who use drugs and/or people who are homeless](#)
- Those at risk and their contacts can be referred to their local public health office for vaccination. For Public Health Office phone numbers and locations please visit: <https://nmhealth.org/location/public/>